

i-Tree

Ecosystem Analysis Prince William County



Urban Forest Effects and Values
March 2021

Summary

Understanding an urban forest's structure, function and value can promote management decisions that will improve human health and environmental quality. An assessment of the vegetation structure, function, and value of the Prince William County urban forest was conducted during 2019 and 2020. Data from 207 field plots located throughout Prince William County were analyzed using the i-Tree Eco model developed by the U.S. Forest Service, Northern Research Station.

- Number of trees: 51,190,000
- Tree Cover: 47.1 %
- Most common species of trees: American beech, Eastern red cedar, Virginia pine
- Percentage of trees less than 6" (15.2 cm) diameter: 74.3%
- Pollution Removal: 5.644 thousand tons/year (\$2.32 billion/year)
- Carbon Storage: 5.451 million tons (\$930 million)
- Carbon Sequestration: 150.9 thousand tons (\$25.7 million/year)
- Oxygen Production: 180.9 thousand tons/year
- Avoided Runoff: 547.7 million cubic feet/year (\$36.6 million/year)
- Building energy savings: \$8,370,000/year
- Carbon Avoided: 12.36 thousand tons/year (\$2110000/year)
- Structural values: \$29 billion

Ton: short ton (U.S.) (2,000 lbs)

Monetary values \$ are reported in US Dollars throughout the report except where noted.

Pollution removal and avoided runoff estimates are reported for trees and shrubs. All other ecosystem service estimates are reported for trees.

For an overview of i-Tree Eco methodology, see Appendix I. Data collection quality is determined by the local data collectors, over which i-Tree has no control. Additionally, some of the plot and tree information may not have been collected, so not all of the analyses may have been conducted for this report.

Table of Contents

| | |
|--|----|
| Summary | 2 |
| I. Tree Characteristics of the Urban Forest | 4 |
| II. Urban Forest Cover and Leaf Area | 7 |
| III. Air Pollution Removal by Urban Trees | 9 |
| IV. Carbon Storage and Sequestration | 11 |
| V. Oxygen Production | 13 |
| VI. Avoided Runoff | 14 |
| VII. Trees and Building Energy Use | 15 |
| VIII. Structural and Functional Values | 16 |
| IX. Potential Pest Impacts | 17 |
| Appendix I. i-Tree Eco Model and Field Measurements | 19 |
| Appendix II. Relative Tree Effects | 23 |
| Appendix III. Comparison of Urban Forests | 24 |
| Appendix IV. General Recommendations for Air Quality Improvement | 25 |
| Appendix V. Invasive Species of the Urban Forest | 26 |
| Appendix VI. Potential Risk of Pests | 27 |
| References | 32 |

I. Tree Characteristics of the Urban Forest

The urban forest of Prince William County has an estimated 51,190,000 trees with a tree cover of 47.1 percent. The three most common species are American beech (11.8 percent), Eastern red cedar (10.4 percent), and Virginia pine (8.2 percent).




Figure 1. Tree species composition in Prince William County

The overall tree density in Prince William County is 256 trees/acre (see Appendix III for comparable values from other cities). For stratified projects, the highest tree densities in Prince William County occur in ParkOpenLand followed by Commercial and AgricultureED.




Figure 2. Number of trees/ac in Prince William County by stratum




Figure 3. Percent of tree population by diameter class (DBH - stem diameter at 4.5 feet)

Urban forests are composed of a mix of native and exotic tree species. Thus, urban forests often have a tree diversity that is higher than surrounding native landscapes. Increased tree diversity can minimize the overall impact or destruction by a species-specific insect or disease, but it can also pose a risk to native plants if some of the exotic species are invasive plants that can potentially out-compete and displace native species. In Prince William County, about 95 percent of the trees are species native to North America, while 95 percent are native to Virginia. Species exotic to North America make up 5 percent of the population. Most exotic tree species have an origin from Asia (3 percent of the species).




Figure 4. Percent of live tree population by area of native origin, Prince William County

The plus sign (+) indicates the tree species is native to another continent other than the ones listed in the grouping.

Invasive plant species are often characterized by their vigor, ability to adapt, reproductive capacity, and general lack of natural enemies. These abilities enable them to displace native plants and make them a threat to natural areas. Eight of the 109 tree species in Prince William County are identified as invasive on the state invasive species list (Virginia Native Plant Society and Department of Conservation and Recreation 2009). These invasive species comprise 2.5 percent of the tree population though they may only cause a minimal level of impact. The three most common invasive species are Autumn olive (1.1 percent of population), Tree of heaven (0.6 percent), and Persian silk tree (0.5 percent) (see Appendix V for a complete list of invasive species).

II. Urban Forest Cover and Leaf Area

Many tree benefits equate directly to the amount of healthy leaf surface area of the plant. Trees cover about 47 percent of Prince William County and provide 2.524 thousand square miles of leaf area. Total leaf area is greatest in ParkOpenLand followed by AgricultureED and HighResidential.




Figure 5. Leaf area by stratum, Prince William County

In Prince William County, the most dominant species in terms of leaf area are American beech, White oak, and Tulip tree. The 10 species with the greatest importance values are listed in Table 1. Importance values (IV) are calculated as the sum of percent population and percent leaf area. High importance values do not mean that these trees should necessarily be encouraged in the future; rather these species currently dominate the urban forest structure.

Table 1. Most important species in Prince William County

| <i>Species Name</i> | <i>Percent Population</i> | <i>Percent Leaf Area</i> | <i>IV</i> |
|---------------------|---------------------------|--------------------------|-----------|
| American beech | 11.8 | 16.3 | 28.0 |
| White oak | 4.0 | 14.2 | 18.2 |
| Red maple | 8.1 | 9.2 | 17.4 |
| Tulip tree | 4.3 | 12.9 | 17.2 |
| Eastern red cedar | 10.4 | 3.1 | 13.5 |
| Virginia pine | 8.2 | 2.9 | 11.1 |
| Black tupelo | 8.1 | 2.5 | 10.6 |
| Northern red oak | 1.5 | 5.2 | 6.7 |
| Mockernut hickory | 2.3 | 3.1 | 5.5 |
| American hornbeam | 3.2 | 1.2 | 4.4 |

Common ground cover classes (including cover types beneath trees and shrubs) in Prince William County include duff/mulch, unmaintained grass, buildings, bare soil, water, rock, and other impervious, impervious covers such as tar, and cement, and herbaceous covers such as grass, and herbs (Figure 6). The most dominant ground cover types are Duff/Mulch (33.2 percent) and Grass (22.8 percent).




Figure 6. Percent of land by ground cover classes, Prince William County

III. Air Pollution Removal by Urban Trees

Poor air quality is a common problem in many urban areas. It can lead to decreased human health, damage to landscape materials and ecosystem processes, and reduced visibility. The urban forest can help improve air quality by reducing air temperature, directly removing pollutants from the air, and reducing energy consumption in buildings, which consequently reduces air pollutant emissions from the power sources. Trees also emit volatile organic compounds that can contribute to ozone formation. However, integrative studies have revealed that an increase in tree cover leads to reduced ozone formation (Nowak and Dwyer 2000).

Pollution removal¹ by trees and shrubs in Prince William County was estimated using field data and recent available pollution and weather data available. Pollution removal was greatest for ozone (Figure 7). It is estimated that trees and shrubs remove 5.644 thousand tons of air pollution (ozone (O3), carbon monoxide (CO), nitrogen dioxide (NO2), particulate matter less than 2.5 microns (PM2.5)², and sulfur dioxide (SO2)) per year with an associated value of \$2.32 billion (see Appendix I for more details).




Figure 7. Annual pollution removal (points) and value (bars) by urban trees, Prince William County

¹ Particulate matter less than 10 microns is a significant air pollutant. Given that i-Tree Eco analyzes particulate matter less than 2.5 microns (PM2.5) which is a subset of PM10, PM10 has not been included in this analysis. PM2.5 is generally more relevant in discussions concerning air pollution effects on human health.

² Trees remove PM2.5 when particulate matter is deposited on leaf surfaces. This deposited PM2.5 can be resuspended to the atmosphere or removed during rain events and dissolved or transferred to the soil. This combination of events can lead to positive or negative pollution removal and value depending on various atmospheric factors (see Appendix I for more details).

In 2019, trees in Prince William County emitted an estimated 7.082 thousand tons of volatile organic compounds (VOCs) (1.706 thousand tons of isoprene and 5.376 thousand tons of monoterpenes). Emissions vary among species based on species characteristics (e.g. some genera such as oaks are high isoprene emitters) and amount of leaf biomass. Fifty- two percent of the urban forest's VOC emissions were from White oak and Northern red oak. These VOCs are precursor chemicals to ozone formation.³

General recommendations for improving air quality with trees are given in Appendix VIII.

³ Some economic studies have estimated VOC emission costs. These costs are not included here as there is a tendency to add positive dollar estimates of ozone removal effects with negative dollar values of VOC emission effects to determine whether tree effects are positive or negative in relation to ozone. This combining of dollar values to determine tree effects should not be done, rather estimates of VOC effects on ozone formation (e.g., via photochemical models) should be conducted and directly contrasted with ozone removal by trees (i.e., ozone effects should be directly compared, not dollar estimates). In addition, air temperature reductions by trees have been shown to significantly reduce ozone concentrations (Cardelino and Chameides 1990; Nowak et al 2000), but are not considered in this analysis. Photochemical modeling that integrates tree effects on air temperature, pollution removal, VOC emissions, and emissions from power plants can be used to determine the overall effect of trees on ozone concentrations.

IV. Carbon Storage and Sequestration

Climate change is an issue of global concern. Urban trees can help mitigate climate change by sequestering atmospheric carbon (from carbon dioxide) in tissue and by altering energy use in buildings, and consequently altering carbon dioxide emissions from fossil-fuel based power sources (Abdollahi et al 2000).

Trees reduce the amount of carbon in the atmosphere by sequestering carbon in new growth every year. The amount of carbon annually sequestered is increased with the size and health of the trees. The gross sequestration of Prince William County trees is about 150.9 thousand tons of carbon per year with an associated value of \$25.7 million. Net carbon sequestration in the urban forest is about 67.84 thousand tons. See Appendix I for more details on methods.




Figure 8. Estimated annual gross carbon sequestration (points) and value (bars) for urban tree species with the greatest sequestration, Prince William County

Carbon storage is another way trees can influence global climate change. As a tree grows, it stores more carbon by holding it in its accumulated tissue. As a tree dies and decays, it releases much of the stored carbon back into the atmosphere. Thus, carbon storage is an indication of the amount of carbon that can be released if trees are allowed to die and decompose. Maintaining healthy trees will keep the carbon stored in trees, but tree maintenance can contribute to carbon emissions (Nowak et al 2002c). When a tree dies, using the wood in long-term wood products, to heat buildings, or to produce energy will help reduce carbon emissions from wood decomposition or from fossil-fuel or wood-based power plants.

Trees in Prince William County are estimated to store 5450000 tons of carbon (\$930 million). Of the species sampled, White oak stores the most carbon (approximately 15.8% of the total carbon stored) and Red maple sequesters the most (approximately 14.1% of all sequestered carbon.)




Figure 9. Estimated carbon storage (points) and values (bars) for urban tree species with the greatest storage, Prince William County

V. Oxygen Production

Oxygen production is one of the most commonly cited benefits of urban trees. The net annual oxygen production of a tree is directly related to the amount of carbon sequestered by the tree, which is tied to the accumulation of tree biomass.

Trees in Prince William County are estimated to produce 180.9 thousand tons of oxygen per year.⁴ However, this tree benefit is relatively insignificant because of the large and relatively stable amount of oxygen in the atmosphere and extensive production by aquatic systems. Our atmosphere has an enormous reserve of oxygen. If all fossil fuel reserves, all trees, and all organic matter in soils were burned, atmospheric oxygen would only drop a few percent (Broecker 1970).

Table 2. The top 20 oxygen production species.

| <i>Species</i> | <i>Oxygen (thousand ton)</i> | <i>Net Carbon Sequestration (thousand ton/yr)</i> | <i>Number of Trees</i> | <i>Leaf Area (square mile)</i> |
|-------------------|----------------------------------|---|------------------------|------------------------------------|
| Red maple | 42.83 | 16.06 | 4,148,190 | 233.42 |
| American beech | 30.09 | 11.28 | 6,036,002 | 410.26 |
| Tulip tree | 29.38 | 11.02 | 2,225,470 | 325.13 |
| Black tupelo | 19.01 | 7.13 | 4,154,291 | 62.54 |
| White oak | 16.27 | 6.10 | 2,066,401 | 357.21 |
| Leyland cypress | 11.26 | 4.22 | 162,375 | 9.25 |
| American elm | 10.61 | 3.98 | 1,031,120 | 48.38 |
| Eastern red cedar | 10.48 | 3.93 | 5,348,192 | 77.57 |
| Black cherry | 9.24 | 3.47 | 595,925 | 21.50 |
| Boxelder | 8.28 | 3.10 | 1,174,313 | 24.28 |
| Black walnut | 7.92 | 2.97 | 341,996 | 30.23 |
| Pin oak | 6.76 | 2.53 | 312,139 | 24.88 |
| Sweetgum | 5.53 | 2.07 | 864,429 | 30.36 |
| American sycamore | 4.80 | 1.80 | 282,332 | 31.01 |
| Willow oak | 4.55 | 1.70 | 269,295 | 42.84 |
| Autumn olive | 4.18 | 1.57 | 552,355 | 4.29 |
| Scarlet oak | 3.95 | 1.48 | 199,901 | 33.72 |
| Flowering dogwood | 3.69 | 1.39 | 853,939 | 6.05 |
| Black oak | 3.65 | 1.37 | 213,166 | 25.89 |
| Virginia pine | 3.41 | 1.28 | 4,188,548 | 73.12 |

⁴ A negative estimate, or oxygen deficit, indicates that trees are decomposing faster than they are producing oxygen. This would be the case in an area that has a large proportion of dead trees.

VI. Avoided Runoff

Surface runoff can be a cause for concern in many urban areas as it can contribute pollution to streams, wetlands, rivers, lakes, and oceans. During precipitation events, some portion of the precipitation is intercepted by vegetation (trees and shrubs) while the other portion reaches the ground. The portion of the precipitation that reaches the ground and does not infiltrate into the soil becomes surface runoff (Hirabayashi 2012). In urban areas, the large extent of impervious surfaces increases the amount of surface runoff.

Urban trees and shrubs, however, are beneficial in reducing surface runoff. Trees and shrubs intercept precipitation, while their root systems promote infiltration and storage in the soil. The trees and shrubs of Prince William County help to reduce runoff by an estimated 548 million cubic feet a year with an associated value of \$37 million (see Appendix I for more details). Avoided runoff is estimated based on local weather from the user-designated weather station. In Prince William County, the total annual precipitation in 2016 was 38.9 inches.




Figure 10. Avoided runoff (points) and value (bars) for species with greatest overall impact on runoff, Prince William County

VII. Trees and Building Energy Use

Trees affect energy consumption by shading buildings, providing evaporative cooling, and blocking winter winds. Trees tend to reduce building energy consumption in the summer months and can either increase or decrease building energy use in the winter months, depending on the location of trees around the building. Estimates of tree effects on energy use are based on field measurements of tree distance and direction to space conditioned residential buildings (McPherson and Simpson 1999).

Trees in Prince William County are estimated to reduce energy-related costs from residential buildings by \$8,370,000 annually. Trees also provide an additional \$2,110,000 in value by reducing the amount of carbon released by fossil-fuel based power plants (a reduction of 12400 tons of carbon emissions).

Note: negative numbers indicate that there was not a reduction in carbon emissions and/or value, rather carbon emissions and values increased by the amount shown as a negative value.⁵

Table 3. Annual energy savings due to trees near residential buildings, Prince William County

| | <i>Heating</i> | <i>Cooling</i> | <i>Total</i> |
|-----------------------|----------------|----------------|--------------|
| MBTU ^a | 174,826 | N/A | 174,826 |
| MWH ^b | 2,765 | 51,095 | 53,861 |
| Carbon Avoided (tons) | 4,561 | 7,798 | 12,359 |

^aMBTU - one million British Thermal Units

^bMWH - megawatt-hour

Table 4. Annual savings ^a(\$) in residential energy expenditure during heating and cooling seasons, Prince William County

| | <i>Heating</i> | <i>Cooling</i> | <i>Total</i> |
|-------------------|----------------|----------------|--------------|
| MBTU ^b | 2,358,452 | N/A | 2,358,452 |
| MWH ^c | 308,485 | 5,700,108 | 6,008,593 |
| Carbon Avoided | 777,838 | 1,329,952 | 2,107,789 |

^bBased on the prices of \$111.558333333333 per MWH and \$13.4903092143955 per MBTU (see Appendix I for more details)

^cMBTU - one million British Thermal Units

^cMWH - megawatt-hour

⁵ Trees modify climate, produce shade, and reduce wind speeds. Increased energy use or costs are likely due to these tree-building interactions creating a cooling effect during the winter season. For example, a tree (particularly evergreen species) located on the southern side of a residential building may produce a shading effect that causes increases in heating requirements.

VIII. Structural and Functional Values

Urban forests have a structural value based on the trees themselves (e.g., the cost of having to replace a tree with a similar tree); they also have functional values (either positive or negative) based on the functions the trees perform.

The structural value of an urban forest tends to increase with a rise in the number and size of healthy trees (Nowak et al 2002a). Annual functional values also tend to increase with increased number and size of healthy trees. Through proper management, urban forest values can be increased; however, the values and benefits also can decrease as the amount of healthy tree cover declines.

Urban trees in Prince William County have the following structural values:

- Structural value: \$29 billion
- Carbon storage: \$930 million

Urban trees in Prince William County have the following annual functional values:

- Carbon sequestration: \$25.7 million
- Avoided runoff: \$36.6 million
- Pollution removal: \$2.32 billion
- Energy costs and carbon emission values: \$10.5 million

(Note: negative value indicates increased energy cost and carbon emission value)




Figure 11. Tree species with the greatest structural value, Prince William County

IX. Potential Pest Impacts

Various insects and diseases can infest urban forests, potentially killing trees and reducing the health, structural value and sustainability of the urban forest. As pests tend to have differing tree hosts, the potential damage or risk of each pest will differ among cities. Thirty-six pests were analyzed for their potential impact and compared with pest range maps (Forest Health Technology Enterprise Team 2014) for the conterminous United States to determine their proximity to Prince William County. Ten of the thirty-six pests analyzed are located within the county. For a complete analysis of all pests, see Appendix VII.




Figure 12. Number of trees at risk (points) and associated compensatory value (bars) for most threatening pests located in the county, Prince William County

Butternut canker (BC) (Ostry et al 1996) is caused by a fungus that infects butternut trees. The disease has since caused significant declines in butternut populations in the United States. Potential loss of trees from BC is 0.7 percent (\$393 million in structural value).

The most common hosts of the fungus that cause chestnut blight (CB) (Diller 1965) are American and European chestnut. CB has the potential to affect 0.0 percent of the population (\$0 in structural value).

Dogwood anthracnose (DA) (Mielke and Daughtrey) is a disease that affects dogwood species, specifically flowering and Pacific dogwood. This disease threatens 1.7 percent of the population, which represents a potential loss of \$115 million in structural value.

American elm, one of the most important street trees in the twentieth century, has been devastated by the Dutch elm disease (DED) (Northeastern Area State and Private Forestry 1998). Since first reported in the 1930s, it has killed over 50 percent of the native elm population in the United States. Although some elm species have shown varying degrees of resistance, Prince William County could possibly lose 2.5 percent of its trees to this pest (\$411 million in structural value).

Emerald ash borer (EAB) (Michigan State University 2010) has killed thousands of ash trees in parts of the United States. EAB has the potential to affect 3.9 percent of the population (\$97.5 million in structural value).

The gypsy moth (GM) (Northeastern Area State and Private Forestry 2005) is a defoliator that feeds on many species causing widespread defoliation and tree death if outbreak conditions last several years. This pest threatens 27.3 percent of the population, which represents a potential loss of \$12.8 billion in structural value.

As one of the most damaging pests to eastern hemlock and Carolina hemlock, hemlock woolly adelgid (HWA) (U.S. Forest Service 2005) has played a large role in hemlock mortality in the United States. HWA has the potential to affect 0.1 percent of the population (\$59.6 million in structural value).

The pine shoot beetle (PSB) (Ciesla 2001) is a wood borer that attacks various pine species, though Scotch pine is the preferred host in North America. PSB has the potential to affect 8.7 percent of the population (\$1.78 billion in structural value).

Although the southern pine beetle (SPB) (Clarke and Nowak 2009) will attack most pine species, its preferred hosts are loblolly, Virginia, pond, spruce, shortleaf, and sand pines. This pest threatens 8.9 percent of the population, which represents a potential loss of \$1.84 billion in structural value.

Thousand cankers disease (TCD) (Cranshaw and Tisserat 2009; Seybold et al 2010) is an insect-disease complex that kills several species of walnuts, including black walnut. Potential loss of trees from TCD is 0.7 percent (\$393 million in structural value).

Appendix I. i-Tree Eco Model and Field Measurements

i-Tree Eco is designed to use standardized field data from randomly located plots and local hourly air pollution and meteorological data to quantify urban forest structure and its numerous effects (Nowak and Crane 2000), including:

- Urban forest structure (e.g., species composition, tree health, leaf area, etc.).
- Amount of pollution removed hourly by the urban forest, and its associated percent air quality improvement throughout a year.
- Total carbon stored and net carbon annually sequestered by the urban forest.
- Effects of trees on building energy use and consequent effects on carbon dioxide emissions from power sources.
- Structural value of the forest, as well as the value for air pollution removal and carbon storage and sequestration.
- Potential impact of infestations by pests, such as Asian longhorned beetle, emerald ash borer, gypsy moth, and Dutch elm disease.

Typically, all field data are collected during the leaf-on season to properly assess tree canopies. Typical data collection (actual data collection may vary depending upon the user) includes land use, ground and tree cover, individual tree attributes of species, stem diameter, height, crown width, crown canopy missing and dieback, and distance and direction to residential buildings (Nowak et al 2005; Nowak et al 2008).

During data collection, trees are identified to the most specific taxonomic classification possible. Trees that are not classified to the species level may be classified by genus (e.g., ash) or species groups (e.g., hardwood). In this report, tree species, genera, or species groups are collectively referred to as tree species.

Tree Characteristics:

Leaf area of trees was assessed using measurements of crown dimensions and percentage of crown canopy missing. In the event that these data variables were not collected, they are estimated by the model.

An analysis of invasive species is not available for studies outside of the United States. For the U.S., invasive species are identified using an invasive species list (Virginia Native Plant Society and Department of Conservation and Recreation 2009) for the state in which the urban forest is located. These lists are not exhaustive and they cover invasive species of varying degrees of invasiveness and distribution. In instances where a state did not have an invasive species list, a list was created based on the lists of the adjacent states. Tree species that are identified as invasive by the state invasive species list are cross-referenced with native range data. This helps eliminate species that are on the state invasive species list, but are native to the study area.

Air Pollution Removal:

Pollution removal is calculated for ozone, sulfur dioxide, nitrogen dioxide, carbon monoxide and particulate matter less than 2.5 microns. Particulate matter less than 10 microns (PM10) is another significant air pollutant. Given that i-Tree Eco analyzes particulate matter less than 2.5 microns (PM2.5) which is a subset of PM10, PM10 has not been included in this analysis. PM2.5 is generally more relevant in discussions concerning air pollution effects on human health.

Air pollution removal estimates are derived from calculated hourly tree-canopy resistances for ozone, and sulfur and nitrogen dioxides based on a hybrid of big-leaf and multi-layer canopy deposition models (Baldocchi 1988; Baldocchi et al 1987). As the removal of carbon monoxide and particulate matter by vegetation is not directly related to transpiration, removal rates (deposition velocities) for these pollutants were based on average measured values from the literature (Bidwell and Fraser 1972; Lovett 1994) that were adjusted depending on leaf phenology and leaf area.

Particulate removal incorporated a 50 percent resuspension rate of particles back to the atmosphere (Zinke 1967). Recent updates (2011) to air quality modeling are based on improved leaf area index simulations, weather and pollution processing and interpolation, and updated pollutant monetary values (Hirabayashi et al 2011; Hirabayashi et al 2012; Hirabayashi 2011).

Trees remove PM_{2.5} when particulate matter is deposited on leaf surfaces (Nowak et al 2013). This deposited PM_{2.5} can be resuspended to the atmosphere or removed during rain events and dissolved or transferred to the soil. This combination of events can lead to positive or negative pollution removal and value depending on various atmospheric factors. Generally, PM_{2.5} removal is positive with positive benefits. However, there are some cases when net removal is negative or resuspended particles lead to increased pollution concentrations and negative values. During some months (e.g., with no rain), trees resuspend more particles than they remove. Resuspension can also lead to increased overall PM_{2.5} concentrations if the boundary layer conditions are lower during net resuspension periods than during net removal periods. Since the pollution removal value is based on the change in pollution concentration, it is possible to have situations when trees remove PM_{2.5} but increase concentrations and thus have negative values during periods of positive overall removal. These events are not common, but can happen.

For reports in the United States, default air pollution removal value is calculated based on local incidence of adverse health effects and national median externality costs. The number of adverse health effects and associated economic value is calculated for ozone, sulfur dioxide, nitrogen dioxide, and particulate matter less than 2.5 microns using data from the U.S. Environmental Protection Agency's Environmental Benefits Mapping and Analysis Program (BenMAP) (Nowak et al 2014). The model uses a damage-function approach that is based on the local change in pollution concentration and population. National median externality costs were used to calculate the value of carbon monoxide removal (Murray et al 1994).

For international reports, user-defined local pollution values are used. For international reports that do not have local values, estimates are based on either European median externality values (van Essen et al 2011) or BenMAP regression equations (Nowak et al 2014) that incorporate user-defined population estimates. Values are then converted to local currency with user-defined exchange rates.

For this analysis, pollution removal value is calculated based on the prices of \$1,327 per ton (carbon monoxide), \$125,616 per ton (ozone), \$37,097 per ton (nitrogen dioxide), \$13,957 per ton (sulfur dioxide), \$5,718,057 per ton (particulate matter less than 2.5 microns).

Carbon Storage and Sequestration:

Carbon storage is the amount of carbon bound up in the above-ground and below-ground parts of woody vegetation. To calculate current carbon storage, biomass for each tree was calculated using equations from the literature and measured tree data. Open-grown, maintained trees tend to have less biomass than predicted by forest-derived biomass equations (Nowak 1994). To adjust for this difference, biomass results for open-grown urban trees were multiplied by 0.8. No adjustment was made for trees found in natural stand conditions. Tree dry-weight biomass was converted to stored carbon by multiplying by 0.5.

Carbon sequestration is the removal of carbon dioxide from the air by plants. To estimate the gross amount of carbon sequestered annually, average diameter growth from the appropriate genera and diameter class and tree condition was added to the existing tree diameter (year x) to estimate tree diameter and carbon storage in year x+1.

Carbon storage and carbon sequestration values are based on estimated or customized local carbon values. For international reports that do not have local values, estimates are based on the carbon value for the United States (U.S. Environmental Protection Agency 2015, Interagency Working Group on Social Cost of Carbon 2015) and converted to local currency with user-defined exchange rates.

For this analysis, carbon storage and carbon sequestration values are calculated based on \$171 per ton.

Oxygen Production:

The amount of oxygen produced is estimated from carbon sequestration based on atomic weights: net O₂ release (kg/yr) = net C sequestration (kg/yr) × 32/12. To estimate the net carbon sequestration rate, the amount of carbon sequestered as a result of tree growth is reduced by the amount lost resulting from tree mortality. Thus, net carbon sequestration and net annual oxygen production of the urban forest account for decomposition (Nowak et al 2007). For complete inventory projects, oxygen production is estimated from gross carbon sequestration and does not account for decomposition.

Avoided Runoff:

Annual avoided surface runoff is calculated based on rainfall interception by vegetation, specifically the difference between annual runoff with and without vegetation. Although tree leaves, branches, and bark may intercept precipitation and thus mitigate surface runoff, only the precipitation intercepted by leaves is accounted for in this analysis.

The value of avoided runoff is based on estimated or user-defined local values. For international reports that do not have local values, the national average value for the United States is utilized and converted to local currency with user-defined exchange rates. The U.S. value of avoided runoff is based on the U.S. Forest Service's Community Tree Guide Series (McPherson et al 1999; 2000; 2001; 2002; 2003; 2004; 2006a; 2006b; 2006c; 2007; 2010; Peper et al 2009; 2010; Vargas et al 2007a; 2007b; 2008).

For this analysis, avoided runoff value is calculated based on the price of \$0.07 per ft³.

Building Energy Use:

If appropriate field data were collected, seasonal effects of trees on residential building energy use were calculated based on procedures described in the literature (McPherson and Simpson 1999) using distance and direction of trees from residential structures, tree height and tree condition data. To calculate the monetary value of energy savings, local or custom prices per MWH or MBTU are utilized.

For this analysis, energy saving value is calculated based on the prices of \$111.56 per MWH and \$13.49 per MBTU.

Structural Values:

Structural value is the value of a tree based on the physical resource itself (e.g., the cost of having to replace a tree with a similar tree). Structural values were based on valuation procedures of the Council of Tree and Landscape Appraisers, which uses tree species, diameter, condition, and location information (Nowak et al 2002a; 2002b). Structural value may not be included for international projects if there is insufficient local data to complete the valuation procedures.

Potential Pest Impacts:

The complete potential pest risk analysis is not available for studies outside of the United States. The number of trees at risk to the pests analyzed is reported, though the list of pests is based on known insects and disease in the United States.

For the U.S., potential pest risk is based on pest range maps and the known pest host species that are likely to

experience mortality. Pest range maps for 2012 from the Forest Health Technology Enterprise Team (FHTET) (Forest Health Technology Enterprise Team 2014) were used to determine the proximity of each pest to the county in which the urban forest is located. For the county, it was established whether the insect/disease occurs within the county, is within 250 miles of the county edge, is between 250 and 750 miles away, or is greater than 750 miles away. FHTET did not have pest range maps for Dutch elm disease and chestnut blight. The range of these pests was based on known occurrence and the host range, respectively (Eastern Forest Environmental Threat Assessment Center; Worrall 2007).

Relative Tree Effects:

The relative value of tree benefits reported in Appendix II is calculated to show what carbon storage and sequestration, and air pollutant removal equate to in amounts of municipal carbon emissions, passenger automobile emissions, and house emissions.

Municipal carbon emissions are based on 2010 U.S. per capita carbon emissions (Carbon Dioxide Information Analysis Center 2010). Per capita emissions were multiplied by city population to estimate total city carbon emissions.

Light duty vehicle emission rates (g/mi) for CO, NO_x, VOCs, PM₁₀, SO₂ for 2010 (Bureau of Transportation Statistics 2010; Heirigs et al 2004), PM_{2.5} for 2011-2015 (California Air Resources Board 2013), and CO₂ for 2011 (U.S. Environmental Protection Agency 2010) were multiplied by average miles driven per vehicle in 2011 (Federal Highway Administration 2013) to determine average emissions per vehicle.

Household emissions are based on average electricity kWh usage, natural gas Btu usage, fuel oil Btu usage, kerosene Btu usage, LPG Btu usage, and wood Btu usage per household in 2009 (Energy Information Administration 2013; Energy Information Administration 2014)

- CO₂, SO₂, and NO_x power plant emission per kWh are from Leonardo Academy 2011. CO emission per kWh assumes 1/3 of one percent of C emissions is CO based on Energy Information Administration 1994. PM₁₀ emission per kWh from Layton 2004.
- CO₂, NO_x, SO₂, and CO emission per Btu for natural gas, propane and butane (average used to represent LPG), Fuel #4 and #6 (average used to represent fuel oil and kerosene) from Leonardo Academy 2011.
- CO₂ emissions per Btu of wood from Energy Information Administration 2014.
- CO, NO_x and SO_x emission per Btu based on total emissions and wood burning (tons) from (British Columbia Ministry 2005; Georgia Forestry Commission 2009).

Appendix II. Relative Tree Effects

The urban forest in Prince William County provides benefits that include carbon storage and sequestration, and air pollutant removal. To estimate the relative value of these benefits, tree benefits were compared to estimates of average municipal carbon emissions, average passenger automobile emissions, and average household emissions. See Appendix I for methodology.

Carbon storage is equivalent to:

- Amount of carbon emitted in Prince William County in 801 days
- Annual carbon (C) emissions from 3,860,000 automobiles
- Annual C emissions from 1,580,000 single-family houses

Carbon monoxide removal is equivalent to:

- Annual carbon monoxide emissions from 785 automobiles
- Annual carbon monoxide emissions from 2,160 single-family houses

Nitrogen dioxide removal is equivalent to:

- Annual nitrogen dioxide emissions from 172,000 automobiles
- Annual nitrogen dioxide emissions from 77,500 single-family houses

Sulfur dioxide removal is equivalent to:

- Annual sulfur dioxide emissions from 543,000 automobiles
- Annual sulfur dioxide emissions from 1,430 single-family houses

Annual carbon sequestration is equivalent to:

- Amount of carbon emitted in Prince William County in 22.0 days
- Annual C emissions from 106,800 automobiles
- Annual C emissions from 43,800 single-family houses

Appendix III. Comparison of Urban Forests

A common question asked is, "How does this city compare to other cities?" Although comparison among cities should be made with caution as there are many attributes of a city that affect urban forest structure and functions, summary data are provided from other cities analyzed using the i-Tree Eco model.

I. City totals for trees

| City | % Tree Cover | Number of Trees | Carbon Storage (tons) | Carbon Sequestration (tons/yr) | Pollution Removal (tons/yr) |
|-----------------------|--------------|-----------------|--------------------------|-----------------------------------|--------------------------------|
| Toronto, ON, Canada | 26.6 | 10,220,000 | 1,221,000 | 51,500 | 2,099 |
| Atlanta, GA | 36.7 | 9,415,000 | 1,344,000 | 46,400 | 1,663 |
| Los Angeles, CA | 11.1 | 5,993,000 | 1,269,000 | 77,000 | 1,975 |
| New York, NY | 20.9 | 5,212,000 | 1,350,000 | 42,300 | 1,676 |
| London, ON, Canada | 24.7 | 4,376,000 | 396,000 | 13,700 | 408 |
| Chicago, IL | 17.2 | 3,585,000 | 716,000 | 25,200 | 888 |
| Phoenix, AZ | 9.0 | 3,166,000 | 315,000 | 32,800 | 563 |
| Baltimore, MD | 21.0 | 2,479,000 | 570,000 | 18,400 | 430 |
| Philadelphia, PA | 15.7 | 2,113,000 | 530,000 | 16,100 | 575 |
| Washington, DC | 28.6 | 1,928,000 | 525,000 | 16,200 | 418 |
| Oakville, ON , Canada | 29.1 | 1,908,000 | 147,000 | 6,600 | 190 |
| Albuquerque, NM | 14.3 | 1,846,000 | 332,000 | 10,600 | 248 |
| Boston, MA | 22.3 | 1,183,000 | 319,000 | 10,500 | 283 |
| Syracuse, NY | 26.9 | 1,088,000 | 183,000 | 5,900 | 109 |
| Woodbridge, NJ | 29.5 | 986,000 | 160,000 | 5,600 | 210 |
| Minneapolis, MN | 26.4 | 979,000 | 250,000 | 8,900 | 305 |
| San Francisco, CA | 11.9 | 668,000 | 194,000 | 5,100 | 141 |
| Morgantown, WV | 35.5 | 658,000 | 93,000 | 2,900 | 72 |
| Moorestown, NJ | 28.0 | 583,000 | 117,000 | 3,800 | 118 |
| Hartford, CT | 25.9 | 568,000 | 143,000 | 4,300 | 58 |
| Jersey City, NJ | 11.5 | 136,000 | 21,000 | 890 | 41 |
| Casper, WY | 8.9 | 123,000 | 37,000 | 1,200 | 37 |
| Freehold, NJ | 34.4 | 48,000 | 20,000 | 540 | 22 |

II. Totals per acre of land area

| City | Number of Trees/ac | Carbon Storage (tons/ac) | Carbon Sequestration (tons/ac/yr) | Pollution Removal (lb/ac/yr) |
|-----------------------|--------------------|-----------------------------|--------------------------------------|---------------------------------|
| Toronto, ON, Canada | 64.9 | 7.8 | 0.33 | 26.7 |
| Atlanta, GA | 111.6 | 15.9 | 0.55 | 39.4 |
| Los Angeles, CA | 19.6 | 4.2 | 0.16 | 13.1 |
| New York, NY | 26.4 | 6.8 | 0.21 | 17.0 |
| London, ON, Canada | 75.1 | 6.8 | 0.24 | 14.0 |
| Chicago, IL | 24.2 | 4.8 | 0.17 | 12.0 |
| Phoenix, AZ | 12.9 | 1.3 | 0.13 | 4.6 |
| Baltimore, MD | 48.0 | 11.1 | 0.36 | 16.6 |
| Philadelphia, PA | 25.1 | 6.3 | 0.19 | 13.6 |
| Washington, DC | 49.0 | 13.3 | 0.41 | 21.2 |
| Oakville, ON , Canada | 78.1 | 6.0 | 0.27 | 11.0 |
| Albuquerque, NM | 21.8 | 3.9 | 0.12 | 5.9 |
| Boston, MA | 33.5 | 9.1 | 0.30 | 16.1 |
| Syracuse, NY | 67.7 | 10.3 | 0.34 | 13.6 |
| Woodbridge, NJ | 66.5 | 10.8 | 0.38 | 28.4 |
| Minneapolis, MN | 26.2 | 6.7 | 0.24 | 16.3 |
| San Francisco, CA | 22.5 | 6.6 | 0.17 | 9.5 |
| Morgantown, WV | 119.2 | 16.8 | 0.52 | 26.0 |
| Moorestown, NJ | 62.1 | 12.4 | 0.40 | 25.1 |
| Hartford, CT | 50.4 | 12.7 | 0.38 | 10.2 |
| Jersey City, NJ | 14.4 | 2.2 | 0.09 | 8.6 |
| Casper, WY | 9.1 | 2.8 | 0.09 | 5.5 |
| Freehold, NJ | 38.3 | 16.0 | 0.44 | 35.3 |

Appendix IV. General Recommendations for Air Quality Improvement

Urban vegetation can directly and indirectly affect local and regional air quality by altering the urban atmosphere environment. Four main ways that urban trees affect air quality are (Nowak 1995):

- Temperature reduction and other microclimate effects
- Removal of air pollutants
- Emission of volatile organic compounds (VOC) and tree maintenance emissions
- Energy effects on buildings

The cumulative and interactive effects of trees on climate, pollution removal, and VOC and power plant emissions determine the impact of trees on air pollution. Cumulative studies involving urban tree impacts on ozone have revealed that increased urban canopy cover, particularly with low VOC emitting species, leads to reduced ozone concentrations in cities (Nowak 2000). Local urban management decisions also can help improve air quality.

Urban forest management strategies to help improve air quality include (Nowak 2000):

| <i>Strategy</i> | <i>Result</i> |
|--|--|
| Increase the number of healthy trees | Increase pollution removal |
| Sustain existing tree cover | Maintain pollution removal levels |
| Maximize use of low VOC-emitting trees | Reduces ozone and carbon monoxide formation |
| Sustain large, healthy trees | Large trees have greatest per-tree effects |
| Use long-lived trees | Reduce long-term pollutant emissions from planting and removal |
| Use low maintenance trees | Reduce pollutants emissions from maintenance activities |
| Reduce fossil fuel use in maintaining vegetation | Reduce pollutant emissions |
| Plant trees in energy conserving locations | Reduce pollutant emissions from power plants |
| Plant trees to shade parked cars | Reduce vehicular VOC emissions |
| Supply ample water to vegetation | Enhance pollution removal and temperature reduction |
| Plant trees in polluted or heavily populated areas | Maximizes tree air quality benefits |
| Avoid pollutant-sensitive species | Improve tree health |
| Utilize evergreen trees for particulate matter | Year-round removal of particles |

Appendix V. Invasive Species of the Urban Forest

The following inventoried tree species were listed as invasive on the Virginia invasive species list (Virginia Native Plant Society and Department of Conservation and Recreation 2009):

| Species Name ^a | Number of Trees | % of Trees | Leaf Area (mi ²) | Percent Leaf Area |
|---------------------------|------------------|-------------|---------------------------------|-------------------|
| Autumn olive | 552,355 | 1.1 | 4.3 | 0.2 |
| Tree of heaven | 315,323 | 0.6 | 11.8 | 0.5 |
| Persian silk tree | 274,627 | 0.5 | 1.1 | 0.0 |
| Amur honeysuckle | 73,168 | 0.1 | 0.4 | 0.0 |
| White mulberry | 27,567 | 0.1 | 1.8 | 0.1 |
| Chinese privet | 21,013 | 0.0 | 0.1 | 0.0 |
| Multiflora rose | 10,318 | 0.0 | 0.0 | 0.0 |
| Royal paulownia | 6,866 | 0.0 | 0.3 | 0.0 |
| Total | 1,281,236 | 2.50 | 19.90 | 0.79 |





^aSpecies are determined to be invasive if they are listed on the state's invasive species list

Appendix VI. Potential Risk of Pests

Thirty-six insects and diseases were analyzed to quantify their potential impact on the urban forest. As each insect/disease is likely to attack different host tree species, the implications for {0} will vary. The number of trees at risk reflects only the known host species that are likely to experience mortality.

| Code | Scientific Name | Common Name | Trees at Risk (#) | Value (\$ billions) |
|-------|--|--------------------------------------|----------------------|------------------------|
| AL | <i>Phyllocnistis populiella</i> | Aspen Leafminer | 198,853 | 0.04 |
| ALB | <i>Anoplophora glabripennis</i> | Asian Longhorned Beetle | 8,012,509 | 3.82 |
| BBD | <i>Neonectria faginata</i> | Beech Bark Disease | 6,036,002 | 2.19 |
| BC | <i>Sirococcus clavignenti juglandacearum</i> | Butternut Canker | 341,996 | 0.39 |
| BWA | <i>Adelges piceae</i> | Balsam Woolly Adelgid | 0 | 0.00 |
| CB | <i>Cryphonectria parasitica</i> | Chestnut Blight | 0 | 0.00 |
| DA | <i>Discula destructiva</i> | Dogwood Anthracnose | 853,939 | 0.12 |
| DBSR | <i>Leptographium wageneri</i> var. <i>pseudotsugae</i> | Douglas-fir Black Stain Root Disease | 0 | 0.00 |
| DED | <i>Ophiostoma novo-ulmi</i> | Dutch Elm Disease | 1,299,004 | 0.41 |
| DFB | <i>Dendroctonus pseudotsugae</i> | Douglas-Fir Beetle | 0 | 0.00 |
| EAB | <i>Agrilus planipennis</i> | Emerald Ash Borer | 1,974,588 | 0.10 |
| FE | <i>Scolytus ventralis</i> | Fir Engraver | 0 | 0.00 |
| FR | <i>Cronartium quercuum</i> f. sp. <i>Fusiforme</i> | Fusiform Rust | 126,699 | 0.00 |
| GM | <i>Lymantria dispar</i> | Gypsy Moth | 13,982,705 | 12.79 |
| GSOB | <i>Agrilus auroguttatus</i> | Goldspotted Oak Borer | 0 | 0.00 |
| HWA | <i>Adelges tsugae</i> | Hemlock Woolly Adelgid | 52,014 | 0.06 |
| JPB | <i>Dendroctonus jeffreyi</i> | Jeffrey Pine Beetle | 0 | 0.00 |
| LAT | <i>Choristoneura conflictana</i> | Large Aspen Tortrix | 406,940 | 0.20 |
| LWD | <i>Raffaelea lauricola</i> | Laurel Wilt | 411,980 | 0.02 |
| MPB | <i>Dendroctonus ponderosae</i> | Mountain Pine Beetle | 0 | 0.00 |
| NSE | <i>Ips perturbatus</i> | Northern Spruce Engraver | 27,085 | 0.01 |
| OW | <i>Ceratocystis fagacearum</i> | Oak Wilt | 4,996,447 | 9.91 |
| PBSR | <i>Leptographium wageneri</i> var. <i>ponderosum</i> | Pine Black Stain Root Disease | 0 | 0.00 |
| POCRD | <i>Phytophthora lateralis</i> | Port-Orford-Cedar Root Disease | 20,702 | 0.01 |
| PSB | <i>Tomicus piniperda</i> | Pine Shoot Beetle | 4,461,883 | 1.78 |
| PSHB | <i>Euwallacea nov. sp.</i> | Polyphagous Shot Hole Borer | 1,174,313 | 0.08 |
| SB | <i>Dendroctonus rufipennis</i> | Spruce Beetle | 27,085 | 0.01 |
| SBW | <i>Choristoneura fumiferana</i> | Spruce Budworm | 27,085 | 0.01 |
| SOD | <i>Phytophthora ramorum</i> | Sudden Oak Death | 2,999,869 | 2.92 |
| SPB | <i>Dendroctonus frontalis</i> | Southern Pine Beetle | 4,540,982 | 1.84 |
| SW | <i>Sirex noctilio</i> | Sirex Wood Wasp | 4,461,883 | 1.78 |
| TCD | <i>Geosmithia morbida</i> | Thousand Cankers Disease | 341,996 | 0.39 |
| WM | <i>Operophtera brumata</i> | Winter Moth | 14,395,421 | 13.87 |
| WPB | <i>Dendroctonus brevicomis</i> | Western Pine Beetle | 0 | 0.00 |
| WPBR | <i>Cronartium ribicola</i> | White Pine Blister Rust | 182,795 | 0.36 |
| WSB | <i>Choristoneura occidentalis</i> | Western Spruce Budworm | 27,085 | 0.01 |

In the following graph, the pests are color coded according to the county's proximity to the pest occurrence in the United States. Red indicates that the pest is within the county; orange indicates that the pest is within 250 miles of the county; yellow indicates that the pest is within 750 miles of the county; and green indicates that the pest is outside of these ranges.



Note: points - Number of trees, bars - Structural value

Based on the host tree species for each pest and the current range of the pest (Forest Health Technology Enterprise Team 2014), it is possible to determine what the risk is that each tree species in the urban forest could be attacked by an insect or disease.

| Spp. Risk | Risk Weight | Species Name | AL | ALB | BBD | BC | BWA | CB | DA | DBSR | DED | DFB | EAB | FE | FR | GM | GSOB | HWA | JPB | LAT | LWD | MPB | NSE | OW | PBSR | POCRD | PSB | PSHB | SB | SBW | SOD | SPB | SW | TCD | WM | WPB | WPBR | WSB | | | |
|-----------|-------------|--------------------|--------|--------|-----|--------|-----|----|----|------|-----|-----|-----|----|----|--------|------|-----|-----|--------|--------|-----|-------|----|------|-------|-----|------|--------|--------|-----|-----|----|-----|--------|--------|--------|--------|--------|--------|--|
| 14 | 14 | Black willow | Yellow | Orange | | | | | | | | | | | | Red | | | | Orange | | | | | | | | | | | | | | | Yellow | | | | | | |
| 14 | 14 | Eastern white pine | | | | | | | | | | | | | | | | | | | | | | | | | Red | | | | | | | | | | | | Orange | | |
| 14 | 14 | Loblolly pine | | | | | | | | | | | | | | Orange | | | | | | | | | | | Red | | | | | | | | | | | | | | |
| 12 | 12 | River birch | | Orange | | | | | | | | | | | | Red | | | | | Orange | | | | | | | | | | | | | | | Yellow | | | | | |
| 12 | 12 | White spruce | | | | | | | | | | | | | | | | | | | | | Green | | | | | | Orange | Orange | | | | | | | | Green | | | |
| 11 | 11 | Virginia pine | | | | | | | | | | | | | | | | | | | | | | | | | Red | | | | | | | | | | | | Orange | | |
| 11 | 11 | Shortleaf pine | | | | | | | | | | | | | | | | | | | | | | | | | Red | | | | | | | | | | | | Orange | | |
| 11 | 11 | pine spp | | | | | | | | | | | | | | | | | | | | | | | | | Red | | | | | | | | | | | | Orange | | |
| 10 | 10 | Boxelder | | Orange | | | | | | | | | | | | | Red | | | | | | | | | | | | Green | | | | | | | | Yellow | | | | |
| 10 | 10 | Northern red oak | | | | | | | | | | | | | | | Red | | | | | | | | | | | | | | | | | | | | | Yellow | | | |
| 10 | 10 | Pin oak | | | | | | | | | | | | | | | Red | | | | | | | | | | | | | | | | | | | | | Yellow | | | |
| 10 | 10 | Southern red oak | | | | | | | | | | | | | | | Red | | | | | | | | | | | | | | | | | | | | | Yellow | | | |
| 10 | 10 | oak spp | | | | | | | | | | | | | | Orange | | | | | | | | | | | | | | | | | | | | | | Yellow | | | |
| 9 | 9 | White oak | | | | | | | | | | | | | | | Red | | | | | | | | | | | | | | | | | | | | | Yellow | | | |
| 9 | 9 | American elm | | Orange | | | | | | | | Red | | | | | | | | | | | | | | | | | | | | | | | | | | | Yellow | | |
| 9 | 9 | Chestnut oak | | | | | | | | | | | | | | | Red | | | | | | | | | | | | | | | | | | | | | | Yellow | | |
| 9 | 9 | Green ash | | Orange | | | | | | | | | Red | | | | | | | | | | | | | | | | | | | | | | | | | | Yellow | | |
| 9 | 9 | Willow oak | | | | | | | | | | | | | | | Red | | | | | | | | | | | | | | | | | | | | | | Yellow | | |
| 9 | 9 | Black oak | | | | | | | | | | | | | | | Red | | | | | | | | | | | | | | | | | | | | | | Yellow | | |
| 9 | 9 | Scarlet oak | | | | | | | | | | | | | | | Red | | | | | | | | | | | | | | | | | | | | | | Yellow | | |
| 9 | 9 | Slippery elm | | Orange | | | | | | | | Red | | | | | | | | | | | | | | | | | | | | | | | | | | | Yellow | | |
| 9 | 9 | Blackjack oak | | | | | | | | | | | | | | | Red | | | | | | | | | | | | | | | | | | | | | | Yellow | | |
| 9 | 9 | Bigtooth aspen | | | | | | | | | | | | | | | | | | | Orange | | | | | | | | | | | | | | | | | | Yellow | | |
| 9 | 9 | Post oak | | | | | | | | | | | | | | | Red | | | | | | | | | | | | | | | | | | | | | | Yellow | | |
| 9 | 9 | Swamp chestnut oak | | | | | | | | | | | | | | | Red | | | | | | | | | | | | | | | | | | | | | | Yellow | | |
| 9 | 9 | Shingle oak | | | | | | | | | | | | | | | Red | | | | | | | | | | | | | | | | | | | | | | Yellow | | |
| 9 | 9 | Swamp white oak | | | | | | | | | | | | | | | Red | | | | | | | | | | | | | | | | | | | | | | Yellow | | |
| 8 | 8 | Black walnut | | | | | Red | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | Red | | | |
| 8 | 8 | Eastern hemlock | | | | | | | | | | | | | | | | Red | | | | | | | | | | | | | | | | | | | | | | Red | |
| 8 | 8 | plum spp | Yellow | Orange | | | | | | | | | | | | | | | | | Orange | | | | | | | | | | | | | | | | | | | Yellow | |
| 7 | 7 | American beech | | | | Orange | | | | | | | | | | | Red | | | | | | | | | | | | | | | | | | | | | | | Yellow | |
| 7 | 7 | elm spp | | Orange | | | | | | | | Red | | | | | | | | | | | | | | | | | | | | | | | | | | | | Yellow | |
| 7 | 7 | apple spp | | Orange | | | | | | | | | | | | | Red | | | | | | | | | | | | | | | | | | | | | | | Yellow | |
| 6 | 6 | White ash | | | | | | | | | | | Red | | | | | | | | | | | | | | | | | | | | | | | | | | Yellow | | |
| 5 | 5 | Red maple | | Orange | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | Yellow | | |
| 5 | 5 | Silver maple | | Orange | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | Yellow | | |

Pest Color Codes:

- Red indicates the pest is within the county
- Orange indicates the pest is within 250 miles of the county
- Yellow indicates pest is within 750 miles of the county
- Green indicates pest is outside of these ranges

References

- Abdollahi, K.K.; Ning, Z.H.; Appeaning, A., eds. 2000. Global climate change and the urban forest. Baton Rouge, LA: GCRCC and Franklin Press. 77 p.
- Baldocchi, D. 1988. A multi-layer model for estimating sulfur dioxide deposition to a deciduous oak forest canopy. *Atmospheric Environment*. 22: 869-884.
- Baldocchi, D.D.; Hicks, B.B.; Camara, P. 1987. A canopy stomatal resistance model for gaseous deposition to vegetated surfaces. *Atmospheric Environment*. 21: 91-101.
- Bidwell, R.G.S.; Fraser, D.E. 1972. Carbon monoxide uptake and metabolism by leaves. *Canadian Journal of Botany*. 50: 1435-1439.
- British Columbia Ministry of Water, Land, and Air Protection. 2005. Residential wood burning emissions in British Columbia. British Columbia.
- Broecker, W.S. 1970. Man's oxygen reserve. *Science* 168(3939): 1537-1538.
- Bureau of Transportation Statistics. 2010. Estimated National Average Vehicle Emissions Rates per Vehicle by Vehicle Type using Gasoline and Diesel. Washington, DC: Bureau of Transportation Statistics, U.S. Department of Transportation. Table 4-43.
- California Air Resources Board. 2013. Methods to Find the Cost-Effectiveness of Funding Air Quality Projects. Table 3 Average Auto Emission Factors. CA: California Environmental Protection Agency, Air Resources Board.
- Carbon Dioxide Information Analysis Center. 2010. CO2 Emissions (metric tons per capita). Washington, DC: The World Bank.
- Cardelino, C.A.; Chameides, W.L. 1990. Natural hydrocarbons, urbanization, and urban ozone. *Journal of Geophysical Research*. 95(D9): 13,971-13,979.
- Ciesla, W. M. 2001. *Tomicus piniperda*. North American Forest Commission. Exotic Forest Pest Information System for North America (EXFOR).
- Clarke, S. R.; Nowak, J.T. 2009. Southern Pine Beetle. Forest Insect & Disease Leaflet 49. Washington, DC: U.S. Department of Agriculture, Forest Service. 8 p.
- Cranshaw, W.; Tisserat, N. 2009. Walnut twig beetle and the thousand cankers disease of black walnut. Pest Alert. Ft. Collins, CO: Colorado State University.
- Seybold, S.; Haugen, D.; Graves, A. 2010. Thousand Cankers Disease. Pest Alert. NA-PR-02-10. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northeastern Area State and Private Forestry.
- Diller, J. D. 1965. Chestnut Blight. Forest Pest Leaflet 94. Washington, DC: U. S. Department of Agriculture, Forest Service. 7 p.
- Eastern Forest Environmental Threat Assessment Center. Dutch Elm Disease. <http://threatsummary.forestthreats.org/threats/threatSummaryViewer.cfm?threatID=43>
- Energy Information Administration. 1994. Energy Use and Carbon Emissions: Non-OECD Countries. Washington, DC:

Energy Information Administration, U.S. Department of Energy.

Energy Information Administration. 2013. CE2.1 Fuel consumption totals and averages, U.S. homes. Washington, DC: Energy Information Administration, U.S. Department of Energy.

Energy Information Administration. 2014. CE5.2 Household wood consumption. Washington, DC: Energy Information Administration, U.S. Department of Energy.

Federal Highway Administration. 2013. Highway Statistics 2011. Washington, DC: Federal Highway Administration, U.S. Department of Transportation. Table VM-1.

Forest Health Technology Enterprise Team. 2014. 2012 National Insect & Disease Risk Maps/Data. Fort Collins, CO: U.S. Department of Agriculture, Forest Service. <http://www.fs.fed.us/foresthealth/technology/nidrm2012.shtml>

Georgia Forestry Commission. 2009. Biomass Energy Conversion for Electricity and Pellets Worksheet. Dry Branch, GA: Georgia Forestry Commission.

Heirigs, P.L.; Delaney, S.S.; Dulla, R.G. 2004. Evaluation of MOBILE Models: MOBILE6.1 (PM), MOBILE6.2 (Toxics), and MOBILE6/CNG. Sacramento, CA: National Cooperative Highway Research Program, Transportation Research Board.

Hirabayashi, S. 2011. Urban Forest Effects-Dry Deposition (UFORE-D) Model Enhancements, [http://www.itreetools.org/eco/resources/UFORE-D enhancements.pdf](http://www.itreetools.org/eco/resources/UFORE-D%20enhancements.pdf)

Hirabayashi, S. 2012. i-Tree Eco Precipitation Interception Model Descriptions, http://www.itreetools.org/eco/resources/iTree_Eco_Precipitation_Interception_Model_Descriptions_V1_2.pdf

Hirabayashi, S.; Kroll, C.; Nowak, D. 2011. Component-based development and sensitivity analyses of an air pollutant dry deposition model. *Environmental Modeling and Software*. 26(6): 804-816.

Hirabayashi, S.; Kroll, C.; Nowak, D. 2012. i-Tree Eco Dry Deposition Model Descriptions V 1.0

Interagency Working Group on Social Cost of Carbon, United States Government. 2015. Technical Support Document: Technical Update of the Social Cost of Carbon for Regulatory Impact Analysis Under Executive Order 12866. <http://www.whitehouse.gov/sites/default/files/omb/inforeg/scc-tsd-final-july-2015.pdf>

Layton, M. 2004. 2005 Electricity Environmental Performance Report: Electricity Generation and Air Emissions. CA: California Energy Commission.

Leonardo Academy. 2011. Leonardo Academy's Guide to Calculating Emissions Including Emission Factors and Energy Prices. Madison, WI: Leonardo Academy Inc.

Lovett, G.M. 1994. Atmospheric deposition of nutrients and pollutants in North America: an ecological perspective. *Ecological Applications*. 4: 629-650.

McPherson, E.G.; Maco, S.E.; Simpson, J.R.; Peper, P.J.; Xiao, Q.; VanDerZanden, A.M.; Bell, N. 2002. Western Washington and Oregon Community Tree Guide: Benefits, Costs, and Strategic Planting. International Society of Arboriculture, Pacific Northwest, Silverton, OR.

McPherson, E.G.; Simpson, J.R. 1999. Carbon dioxide reduction through urban forestry: guidelines for professional and volunteer tree planters. Gen. Tech. Rep. PSW-171. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station. 237 p.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Crowell, A.M.N.; Xiao, Q. 2010. Northern California coast community tree guide: benefits, costs, and strategic planting. PSW-GTR-228. Gen. Tech. Rep. PSW-GTR-228. U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, Albany, CA.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Gardner, S.L.; Vargas, K.E.; Maco, S.E.; Xiao, Q. 2006a. Coastal Plain Community Tree Guide: Benefits, Costs, and Strategic Planting PSW-GTR-201. USDA Forest Service, Pacific Southwest Research Station, Albany, CA.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Gardner, S.L.; Vargas, K.E.; Xiao, Q. 2007. Northeast community tree guide: benefits, costs, and strategic planting.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Maco, S.E.; Gardner, S.L.; Cozad, S.K.; Xiao, Q. 2006b. Midwest Community Tree Guide: Benefits, Costs and Strategic Planting PSW-GTR-199. U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, Albany, CA.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Maco, S.E.; Gardner, S.L.; Vargas, K.E.; Xiao, Q. 2006c. Piedmont Community Tree Guide: Benefits, Costs, and Strategic Planting PSW-GTR 200. U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, Albany, CA.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Maco, S.E.; Xiao Q.; Mulrean, E. 2004. Desert Southwest Community Tree Guide: Benefits, Costs and Strategic Planting. Phoenix, AZ: Arizona Community Tree Council, Inc. 81 :81.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Scott, K.I.; Xiao, Q. 2000. Tree Guidelines for Coastal Southern California Communities. Local Government Commission, Sacramento, CA.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Xiao, Q. 1999. Tree Guidelines for San Joaquin Valley Communities. Local Government Commission, Sacramento, CA.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Xiao, Q.; Maco, S.E.; Hoefer, P.J. 2003. Northern Mountain and Prairie Community Tree Guide: Benefits, Costs and Strategic Planting. Center for Urban Forest Research, USDA Forest Service, Pacific Southwest Research Station, Albany, CA.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Xiao, Q.; Pittenger, D.R.; Hodel, D.R. 2001. Tree Guidelines for Inland Empire Communities. Local Government Commission, Sacramento, CA.

Michigan State University. 2010. Emerald ash borer. East Lansing, MI: Michigan State University [and others].

Mielke, M. E.; Daughtrey, M. L. How to Identify and Control Dogwood Anthracnose. NA-GR-18. Broomall, PA: U. S. Department of Agriculture, Forest Service, Northeastern Area and Private Forestry.

Murray, F.J.; Marsh L.; Bradford, P.A. 1994. New York State Energy Plan, vol. II: issue reports. Albany, NY: New York State Energy Office.

National Invasive Species Information Center. 2011. Beltsville, MD: U.S. Department of Agriculture, National Invasive Species Information Center. <http://www.invasivespeciesinfo.gov/plants/main.shtml>

Northeastern Area State and Private Forestry. 1998. How to identify and manage Dutch Elm Disease. NA-PR-07-98. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northeastern Area State and Private Forestry.

Northeastern Area State and Private Forestry. 2005. Gypsy moth digest. Newtown Square, PA: U.S. Department of

Agriculture, Forest Service, Northeastern Area State and Private Forestry.

Nowak, D.J. 1994. Atmospheric carbon dioxide reduction by Chicago's urban forest. In: McPherson, E.G.; Nowak, D.J.; Rowntree, R.A., eds. Chicago's urban forest ecosystem: results of the Chicago Urban Forest Climate Project. Gen. Tech. Rep. NE-186. Radnor, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station: 83-94.

Nowak, D.J. 1995. Trees pollute? A "TREE" explains it all. In: Proceedings of the 7th National Urban Forestry Conference. Washington, DC: American Forests: 28-30.

Nowak, D.J. 2000. The interactions between urban forests and global climate change. In: Abdollahi, K.K.; Ning, Z.H.; Appeaning, A., eds. Global Climate Change and the Urban Forest. Baton Rouge, LA: GCRCC and Franklin Press: 31-44.

Nowak, D.J., Hirabayashi, S., Bodine, A., Greenfield, E. 2014. Tree and forest effects on air quality and human health in the United States. *Environmental Pollution*. 193:119-129.

Nowak, D.J., Hirabayashi, S., Bodine, A., Hoehn, R. 2013. Modeled PM_{2.5} removal by trees in ten U.S. cities and associated health effects. *Environmental Pollution*. 178: 395-402.

Nowak, D.J.; Civerolo, K.L.; Rao, S.T.; Sistla, S.; Luley, C.J.; Crane, D.E. 2000. A modeling study of the impact of urban trees on ozone. *Atmospheric Environment*. 34: 1601-1613.

Nowak, D.J.; Crane, D.E. 2000. The Urban Forest Effects (UFORE) Model: quantifying urban forest structure and functions. In: Hansen, M.; Burk, T., eds. Integrated tools for natural resources inventories in the 21st century. Proceedings of IUFRO conference. Gen. Tech. Rep. NC-212. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Research Station: 714-720.

Nowak, D.J.; Crane, D.E.; Dwyer, J.F. 2002a. Compensatory value of urban trees in the United States. *Journal of Arboriculture*. 28(4): 194 - 199.

Nowak, D.J.; Crane, D.E.; Stevens, J.C.; Hoehn, R.E. 2005. The urban forest effects (UFORE) model: field data collection manual. V1b. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northeastern Research Station, 34 p. http://www.fs.fed.us/ne/syracuse/Tools/downloads/UFORE_Manual.pdf

Nowak, D.J.; Crane, D.E.; Stevens, J.C.; Ibarra, M. 2002b. Brooklyn's urban forest. Gen. Tech. Rep. NE-290. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northeastern Research Station. 107 p.

Nowak, D.J.; Dwyer, J.F. 2000. Understanding the benefits and costs of urban forest ecosystems. In: Kuser, John, ed. Handbook of urban and community forestry in the northeast. New York, NY: Kluwer Academics/Plenum: 11-22.

Nowak, D.J.; Hoehn, R.; Crane, D. 2007. Oxygen production by urban trees in the United States. *Arboriculture & Urban Forestry*. 33(3):220-226.

Nowak, D.J.; Hoehn, R.E.; Crane, D.E.; Stevens, J.C.; Walton, J.T; Bond, J. 2008. A ground-based method of assessing urban forest structure and ecosystem services. *Arboriculture and Urban Forestry*. 34(6): 347-358.

Nowak, D.J.; Stevens, J.C.; Sisinni, S.M.; Luley, C.J. 2002c. Effects of urban tree management and species selection on atmospheric carbon dioxide. *Journal of Arboriculture*. 28(3): 113-122.

Ostry, M.E.; Mielke, M.E.; Anderson, R.L. 1996. How to Identify Butternut Canker and Manage Butternut Trees. U. S. Department of Agriculture, Forest Service, North Central Forest Experiment Station.

Peper, P.J.; McPherson, E.G.; Simpson, J.R.; Albers, S.N.; Xiao, Q. 2010. Central Florida community tree guide: benefits, costs, and strategic planting. Gen. Tech. Rep. PSW-GTR-230. U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, Albany, CA.

Peper, P.J.; McPherson, E.G.; Simpson, J.R.; Vargas, K.E.; Xiao Q. 2009. Lower Midwest community tree guide: benefits, costs, and strategic planting. PSW-GTR-219. Gen. Tech. Rep. PSW-GTR-219. U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, Albany, CA.

U.S. Environmental Protection Agency. 2010. Light-Duty Vehicle Greenhouse Gas Emission Standards and Corporate Average Fuel Economy Standards. Washington, DC: U.S. Environmental Protection Agency. EPA-420-R-10-012a

U.S. Environmental Protection Agency. 2015. The social cost of carbon. <http://www.epa.gov/climatechange/EPAactivities/economics/scc.html>

U.S. Forest Service. 2005. Hemlock Woolly Adelgid. Pest Alert. NA-PR-09-05. Newtown Square, PA: U. S. Department of Agriculture, Forest Service, Northern Area State and Private Forestry.

van Essen, H.; Schroten, A.; Otten, M.; Sutter, D.; Schreyer, C.; Zandonella, R.; Maibach, M.; Doll, C. 2011. External Costs of Transport in Europe. Netherlands: CE Delft. 161 p.

Vargas, K.E.; McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Gardner, S.L.; Xiao, Q. 2007a. Interior West Tree Guide.

Vargas, K.E.; McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Gardner, S.L.; Xiao, Q. 2007b. Temperate Interior West Community Tree Guide: Benefits, Costs, and Strategic Planting.

Vargas, K.E.; McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Gardner, S.L.; Xiao, Q. 2008. Tropical community tree guide: benefits, costs, and strategic planting. PSW-GTR-216. Gen. Tech. Rep. PSW-GTR-216. U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, Albany, CA.

Virginia Native Plant Society; Department of Conservation and Recreation. 2009. Invasive Alien Plant Species of Virginia. Richmond, VA: Department of Conservation and Recreation, Division of Natural Heritage. <http://www.dcr.virginia.gov/natural_heritage/documents/invlist.pdf>

Worrall, J.J. 2007. Chestnut Blight. Forest and Shade Tree Pathology. http://www.forestpathology.org/dis_chestnut.html

Zinke, P.J. 1967. Forest interception studies in the United States. In: Sopper, W.E.; Lull, H.W., eds. Forest Hydrology. Oxford, UK: Pergamon Press: 137-161.